

HEPATITIS A

General Information

What is hepatitis?

“Hepatitis” means inflammation of the liver. The liver is a vital organ that processes nutrients, filters the blood, and fights infections. When the liver is inflamed or damaged, its function can be affected.

Hepatitis is most often caused by a virus. In the United States, the most common types of viral hepatitis are Hepatitis A, Hepatitis B, and Hepatitis C. Heavy alcohol use, toxins, some medications, and certain medical conditions can also cause hepatitis.

What is Hepatitis A?

Hepatitis A is a highly contagious liver infection caused by the Hepatitis A virus. It can range in severity from a mild illness lasting a few weeks to a severe illness lasting several months.

Who is at risk?

Although anyone can get Hepatitis A, some people are at greater risk, such as those who:

- Travel to or live in countries where Hepatitis A is common
- Use recreational drugs, whether injected or not
- Have sexual contact with someone who has Hepatitis A
- Have clotting-factor disorders, such as hemophilia
- Are men who have sexual encounters with other men
- Are household members or caregivers of a person infected with Hepatitis A

How common is Hepatitis A?

Hepatitis A still occurs in the United States, although not as frequently as it once did. Over the last several decades, there has been more than a 90% decrease in Hepatitis A cases. New cases are now estimated to be around 3,000 each year. Many experts believe this decline is a result of the vaccination of children and people at risk for Hepatitis A. Many of the new cases, however, are from American travelers who got infected while traveling to parts of the world where Hepatitis A is common.



Hepatitis A can be prevented with a safe and effective vaccine.

How is Hepatitis A spread?

Hepatitis A is usually spread when a person ingests fecal matter—even in microscopic amounts—from contact with objects, food, or drinks contaminated by feces or stool from an infected person.

Hepatitis A can be spread when:

- An infected person does not wash his/her hands properly after going to the bathroom and then touches objects or food
- A caregiver does not properly wash his or her hands after changing diapers or cleaning up the stool of an infected person
- Someone engages in sexual activities with an infected person

Hepatitis A also can be spread through contaminated food or water. Contamination of food can happen at any point: growing, harvesting, processing, handling, and even after cooking. This most often occurs in countries where Hepatitis A is common.

Continued on next page



What are the symptoms of Hepatitis A?

Not everyone has symptoms. If symptoms develop, they usually appear 2 to 6 weeks after infection and can include:

- Fever
- Vomiting
- Grey-colored stools
- Fatigue
- Abdominal pain
- Joint pain
- Loss of appetite
- Dark urine
- Jaundice
- Nausea

Symptoms are more likely to occur in adults than in children. They usually last less than 2 months, although some people can be ill for as long as 6 months.



People can spread Hepatitis A even if they don't look or feel sick. Many children and some adults have no symptoms.

How is Hepatitis A diagnosed and treated?

A doctor can determine if a person has Hepatitis A by discussing his or her symptoms and taking a blood sample. To treat Hepatitis A, doctors usually recommend rest, adequate nutrition, fluids, and medical monitoring. Some people will need to be hospitalized. It can take a few months before people begin to feel better.

How serious is Hepatitis A?

Most people who get Hepatitis A feel sick for several months, but they usually recover completely and do not have lasting liver damage. Sometimes Hepatitis A can cause liver failure and death, although this is rare and occurs more commonly in people older than 50 and people with other liver diseases.

Can Hepatitis A be prevented?

Yes. The best way to prevent Hepatitis A is by getting vaccinated. Experts recommend the vaccine for all children, and people with certain risk factors and medical conditions. The vaccine is also recommended for travelers to certain international countries, even if travel occurs for short times or on closed resorts. The Hepatitis A vaccine is safe and effective and given as 2 shots, 6 months apart. Both shots are needed for long-term protection. Ask if your health plan will cover travel related vaccines. You can get vaccinated at your doctor's office, as well as travel clinics and other locations. Lower cost vaccination may be available at certain pharmacies and your local health department.

Who should get vaccinated against Hepatitis A?

Vaccination is recommended for certain groups, including:

- All children at age 1 year
- Travelers to countries where Hepatitis A is common
- Family and caregivers of adoptees from countries where Hepatitis A is common
- Men who have sexual encounters with other men
- Users of recreational drugs, whether injected or not
- People with chronic or long-term liver disease, including Hepatitis B or Hepatitis C
- People with clotting-factor disorders

For more information

Talk to your health professional, call your health department, or visit www.cdc.gov/hepatitis or www.cdc.gov/travel.

HEPATITIS B

General Information

What is hepatitis?

"Hepatitis" means inflammation of the liver. The liver is a vital organ that processes nutrients, filters the blood, and fights infections. When the liver is inflamed or damaged, its function can be affected. Heavy alcohol use, toxins, some medications, and certain medical conditions can cause hepatitis. However, hepatitis is most often caused by a virus. In the United States, the most common types of viral hepatitis are Hepatitis A, Hepatitis B, and Hepatitis C.



The only way to know if you have Hepatitis B is to get tested.

What is Hepatitis B?

Hepatitis B can be a serious liver disease that results from infection with the Hepatitis B virus. **Acute Hepatitis B** refers to a short-term infection that occurs within the first 6 months after someone is infected with the virus. The infection can range in severity from a mild illness with few or no symptoms to a serious condition requiring hospitalization. Some people, especially adults, are able to clear, or get rid of, the virus without treatment. People who clear the virus become immune and cannot get infected with the Hepatitis B virus again.

Chronic Hepatitis B refers to a lifelong infection with the Hepatitis B virus. The likelihood that a person develops a chronic infection depends on the age at which someone becomes infected. Up to 90% of infants infected with the Hepatitis B virus will develop a chronic infection. In contrast, about 5% of adults will develop chronic Hepatitis B. Over time, chronic Hepatitis B can cause serious health problems, including liver damage, cirrhosis, liver cancer, and even death.

How is Hepatitis B spread?

The Hepatitis B virus is spread when blood, semen, or other body fluids from an infected person enters the body of someone who is not infected. The virus can be spread through:

- **Sex with an infected person.** Among adults, Hepatitis B is often spread through sexual contact.
- **Injection drug use.** Sharing needles, syringes, and any other equipment to inject drugs with someone infected with Hepatitis B can spread the virus.
- **Outbreaks.** While uncommon, poor infection control has resulted in outbreaks of Hepatitis B in healthcare settings.
- **Birth.** Hepatitis B can be passed from an infected mother to her baby at birth. Worldwide, most people with Hepatitis B were infected with the virus as an infant.

Hepatitis B is **not** spread through breastfeeding, sharing eating utensils, hugging, kissing, holding hands, coughing, or sneezing. Unlike some forms of hepatitis, Hepatitis B is also not spread by contaminated food or water.

What are the symptoms of Hepatitis B?

Many people with Hepatitis B do not have symptoms and do not know they are infected. If symptoms occur, they can include: fever, feeling tired, not wanting to eat, upset stomach, throwing up, dark urine, grey-colored stool, joint pain, and yellow skin and eyes.

When do symptoms occur?

If symptoms occur with an acute infection, they usually appear within 3 months of exposure and can last up to 6 months. If symptoms occur with chronic Hepatitis B, they can take years to develop and can be a sign of advanced liver disease.

Continued on next page



How would you know if you have Hepatitis B?

The only way to know if you have Hepatitis B is to get tested. Blood tests can determine if a person has been infected and cleared the virus, is currently infected, or has never been infected.

Who should get tested for Hepatitis B and why?

CDC develops recommendations for testing based upon a variety of different factors. Here is a list of people who should get tested. The results will help determine the next best steps for vaccination or medical care.

All pregnant women are routinely tested for Hepatitis B. If a woman has Hepatitis B, timely vaccination can help prevent the spread of the virus to her baby.

Household and sexual contacts of people with Hepatitis B are at risk for getting Hepatitis B. Those who have never had Hepatitis B can benefit from vaccination.

People born in certain parts of the world that have increased rates of Hepatitis B. Testing helps identify those who are infected so that they can receive timely medical care.

People with certain medical conditions should be tested, and get vaccinated if needed. This includes people with HIV infection, people who receive chemotherapy and people on hemodialysis.

People who inject drugs are at increased risk for Hepatitis B but testing can tell if someone is infected or could benefit from vaccination to prevent getting infected with the virus.

Men who have sex with men have higher rates of Hepatitis B. Testing can identify unknown infections or let a person know that they can benefit from vaccination.

How is Hepatitis B treated?

For those with acute Hepatitis B, doctors usually recommend rest, adequate nutrition, fluids, and close medical monitoring. Some people may need to be hospitalized. People living with chronic Hepatitis B should be evaluated for liver problems and monitored on a regular basis. Treatments are available that can slow down or prevent the effects of liver disease.

Can Hepatitis B be prevented?

Yes. The best way to prevent Hepatitis B is by getting vaccinated. The Hepatitis B vaccine is typically given as a series of 3 shots over a period of 6 months. The entire series is needed for long-term protection.

Who should get vaccinated against Hepatitis B?

All infants are routinely vaccinated for Hepatitis B at birth, which has led to dramatic declines of new Hepatitis B cases in the US and many parts of the world. The vaccine is also recommended for people living with someone infected with Hepatitis B, travelers to certain countries, and healthcare and public safety workers exposed to blood. People with high-risk sexual behaviors, men who have sex with men, people who inject drugs, and people who have certain medical conditions, including diabetes, should talk to their doctor about getting vaccinated.

For more information

Talk to your doctor, call your health department, or visit www.cdc.gov/hepatitis.